

LETTER • OPEN ACCESS

Temperature and precipitation extremes under current, 1.5 °C and 2.0 °C global warming above pre-industrial levels over Botswana, and implications for climate change vulnerability

To cite this article: Tiro Nkemelang *et al* 2018 *Environ. Res. Lett.* **13** 065016

View the [article online](#) for updates and enhancements.

Related content

- [On the need for regional climate information over Africa under varying levels of global warming](#)
C J Lennard, G Nikulin, A Dosio *et al.*
- [The southern African climate under 1.5 °C and 2 °C of global warming as simulated by CORDEX regional climate models](#)
G Maúre, I Pinto, M Ndebele-Murisa *et al.*
- [Changes in climate extremes over West and Central Africa at 1.5 °C and 2 °C global warming](#)
Arona Diedhiou, Adeline Bichet, Richard Wartenburger *et al.*

Recent citations

- [Regional climate projections for impact assessment studies in East Africa](#)
Solomon H Gebrechorkos *et al*

Environmental Research Letters



LETTER

OPEN ACCESS

RECEIVED
31 October 2017

REVISED
2 May 2018

ACCEPTED FOR PUBLICATION
8 May 2018

PUBLISHED
14 June 2018

Original content from
this work may be used
under the terms of the
[Creative Commons
Attribution 3.0 licence](#).

Any further distribution
of this work must
maintain attribution to
the author(s) and the
title of the work, journal
citation and DOI.



Temperature and precipitation extremes under current, 1.5 °C and 2.0 °C global warming above pre-industrial levels over Botswana, and implications for climate change vulnerability

Tiro Nkemelang^{1,2,7} , Mark New^{2,3,4}  and Modathir Zaroug^{5,2,6}

¹ Botswana Institute for Technology Research and Innovation, Gaborone, Botswana

² African Climate and Development Initiative, University of Cape Town, Cape Town, South Africa

³ Department of Environmental and Geographical Science, University of Cape Town, Cape Town, South Africa

⁴ School of International Development, University of East Anglia, Norwich, United Kingdom

⁵ Nile Basin Initiative, Kampala, Uganda

⁶ Climate System Analysis Group, University of Cape Town, South Africa

⁷ Author to whom any correspondence should be addressed.

E-mail: tnkemelang@bitri.co.bw

Keywords: climate extremes, Botswana, 1.5 degrees, climate change, Africa, climate impacts

Supplementary material for this article is available [online](#)

Abstract

Climate extremes are widely projected to become more severe as the global climate continues to warm due to anthropogenic greenhouse gas emissions. These extremes often cause the most severe impacts on society. Therefore, the extent to which the extremes might change at regional level as the global climate warms from current levels to proposed policy targets of 1.5 and 2.0 °C above preindustrial levels need to be understood to allow for better preparedness and informed policy formulation. This paper analysed projected changes in temperature and precipitation extremes at 1.0, 1.5 and 2.0 °C warming over Botswana, a country highly vulnerable to the impacts of climate change. Projected changes in temperature extremes are significantly different from each other at the three levels of global warming, across three main climatic zones in the country. Specifically, at 2.0 °C global warming relative to preindustrial, for the ensemble median: (a) country average warm spell duration index increases by 80, 65, 62 days per year across different climatic zones, approximately three (and two) times the change at 1.0 (1.5) °C; (b) cold night (TN10P) and cold day (TX10P) frequencies decrease by 12 and 9 days per year across all regions, respectively, while hot nights (TN90P) and hot days (TX90P) both increase by 8–9 days across all regions. Projected changes in drought-related indices are also distinct at different warming levels. Specifically: (a) projected mean annual precipitation decreases across the country by 5%–12% at 2 °C, 3%–8% at 1.5 °C and 2%–7% at 1.0 °C; (b) dry spell length (ALTCDD) increases by 15–19 days across the three climatic zones at 2.0 °C, about three (and two) times as much as the increase at 1.0 (1.5) °C. Ensemble mean projections indicate increases in heavy rainfall indices, but the inter-model spread is large, with no consistent direction of change, and so changes are not statistically significant. The implications of these changes in extreme temperature and precipitation for key socio-economic sectors are explored, and reveal progressively severe impacts, and consequent adaptation challenges for Botswana as the global climate warms from its present temperature of 1.0 °C above preindustrial levels to 1.5 °C, and then 2.0 °C.

1. Introduction

Some of the harshest impacts of climate change are likely to occur through more extreme climate

and weather events (Trenberth 2012). Sub-Saharan Africa is among the most vulnerable of regions (Boko *et al* 2008, Niang *et al* 2014), with 34 of the top 50 most climate-vulnerable countries located in the

continent (ND-GAIN 2016). Therefore, climate change and extremes will continue to pose significant social and economic pressures on livelihoods within the continent under a warming global climate (King *et al* 2015, ASC 2016). The frequency and intensity of extreme weather events is set to change, often increasing, across most regions thereby increasing vulnerability of those already exposed and sensitive especially rural communities (McElroy and Baker 2012, Adams *et al* 2013). These extreme events, including heat waves (Ceccherini *et al* 2017), heavy precipitation events causing floods (Cook *et al* 2004, Reason 2007, Manhique *et al* 2015, Moyo and Nangombe 2015) as well as droughts (Richard *et al* 2001, Rouault and Richard 2005) are already common drivers of vulnerability in southern Africa.

The 2015 Paris climate change accord, drawing on growing evidence from successive IPCC reports, agreed to work to keeping global mean surface temperature (GMST) warming due to greenhouse gas and other emissions to below 2.0 °C, with the ambition to keep temperature change below 1.5 °C of pre-industrial temperature (UNFCCC 2015). Based on these two temperature warmings, various studies have been conducted on shifts in climates of different localities across the globe, assessing the extent to which an additional 0.5 °C temperature rise would influence both the mean and the climate extremes (e.g. James *et al* 2017, King *et al* 2017, Sanderson *et al* 2017, Wang *et al* 2017). With some studies proposing that the 1.5 °C warming level could be crossed as early as 2026 (Henley and King 2017), the need for early action becomes even more critical. This therefore calls for more detailed analysis of the potential regional and local changes in extremes that might accompany these levels of global temperature increase, especially in the most vulnerable regions of the world.

Previous work on assessing observed and projected trends in climatic extremes have largely been based on a set of 27 temperature and precipitation indices, devised by the Expert Team on Climate Change Detection and Indices (ETCCDI) (Zhang *et al* 2011). Several studies have shown changes in these indices over the last several decades across the globe, based on observed data (e.g. Frich *et al* 2002, Alexander *et al* 2006, Manton 2010) as well reanalysis datasets (e.g. Sillmann *et al* 2013a). These studies have shown a general trend towards more extreme weather under climate change. For future climate change, a number of studies have looked at the projected changes in both extremes of temperature (e.g. Sillmann *et al* 2013b, Lewis and King 2017) and precipitation (Sillmann *et al* 2013b) globally. Most of these studies looked at these changes for a particular period in the 21st century following a particular Representative Concentration Pathway (RCP) (Fischer *et al* 2013, Lewis and King 2017) or emissions scenario (Kharin *et al* 2007).

At regional scales, a number of studies have been carried out to assess both the historical and pro-

jected future trends in temperature and precipitation extremes (e.g. You *et al* 2011, Monier and Gao 2015, Dosio 2016, Razavi *et al* 2016). For Southern Africa, New *et al* (2006) investigated the historical context of extremes from observations, while Shongwe *et al* (2009) and Pinto *et al* (2016) looked at the historical and future projection of precipitation extremes from reanalysis and downscaled climate model outputs. There has so far been little research on the relative impact of global warming of 1.5 °C and 2.0 °C on climate extremes at the national or sub-national scale.

Botswana, though highly exposed and vulnerable to the impacts of climate change, is under-researched on physical climate change, especially on the evolution of climate extremes. A recent study ranked Botswana among the top few countries in Africa in terms of national mean temperature and precipitation changes as global temperature rises to 1.5 °C and 2.0 °C (Zaroug *et al* 2018). This potential vulnerability is reflected in policy and practice in Botswana: Hambira and Saarinen (2015) note the widespread agreement among policy makers on the perceived worsening climate patterns in Botswana; Akinyemi (2017) reports similar perceptions amongst farming communities in Eastern Botswana; in both these cases, there is no quantitative research available to back up these views. This paper therefore aims to quantify the projected changes in climate extremes that can be expected at the global temperature targets agreed in Paris, relative to both pre-industrial and present-day global temperature levels. The paper then discusses the implications of these projected changes for vulnerability of key sectors in Botswana.

2. Data and methodology

2.1. Study area

Figure 1(a)), sharing borders with South Africa to the south, Namibia to the west, Zambia to the north and Zimbabwe to the north east. The subsiding limb of the tropical Hadley circulation defines much of the climate in Botswana, with a semi-permanent high pressure system (the Botswana high) persistent over the region (Driver and Reason 2017). The south-western part of the country is hyper-arid, receiving the lowest rainfall, with aridity then decreasing to North and East of the country (figure 1(b)). Temperatures in Botswana are lowest during the austral winter and highest during the austral summer. In winter, the occasional passing of westerly cold frontal systems can cause minimum temperatures to fall to below freezing resulting in frosts over most parts of the country (Andringa 1984). Summer mean maximum temperatures range between 30.9 °C–33.0 °C across the country, the western parts of the country being hotter (Moses 2017). Occasional heat waves are also experienced in austral summer with temperatures reaching highs of over 42 °C in some places (Moses 2017). Botswana's




Figure 1. (a) Map of Southern Africa with Botswana highlighted (Mapchart.net). (b) Mean annual precipitation (1979–2013) in mm over Botswana derived from the WFDEI dataset and the three regions of homogeneous monthly rainfall in Botswana (Region 1, 2 and 3).

interannual climate variability is mainly driven by the El Niño Southern Oscillation (ENSO), where El Niño events are associated with dry conditions (Nicholson *et al* 2001) and La Niña events are associated with wet years (Mason and Jury 1997). Rain bearing weather systems affecting the country include among others temperate tropical troughs (Williams *et al* 2007) and mesoscale convective systems (Blamey and Reason 2012). These systems are mainly convection driven and associated with the southward movement of the Intertropical Convergence Zone (ITCZ) in the austral summer months (mainly November to March) (Bhalotra 1987). Tropical depressions that form in the Indian Ocean also occasionally penetrate the subcontinent from the east, bringing along heavy precipitation (Reason and Keibel 2004, Reason 2007). Some heavy precipitation events could be a combination of various weather systems such as westerly troughs (cold fronts), tropical lows and ridging anticyclones creating conducive environment for such events (Crimp and Mason 1999). Cut off lows are also a common occurrence over the region, they tend to bring along heavy precipitation episodes that can cause flooding in some places in Southern Africa including Botswana (Singleton and Reason 2007, Favre *et al* 2013, Molekwa 2013). Cut off lows also tend to be associated with extreme rainy days that fall outside of the normal rainy season (Favre *et al* 2013). Dry spells within the rainy season are also a common occurrence in Botswana (Vossen 1990), being more frequent during ENSO warm phases and over the south-western parts of the country (Usman and Reason 2004).

2.2. Data

The ETCCDI temperature and precipitation extreme indices derived from the fifth version of Coupled Model Intercomparison Project (CMIP5) program

participating models were analyzed. A total of 24 CMIP5 Global Climate Model (GCM) outputs developed by Sillmann *et al* (2013a) (table S1 available at stacks.iop.org/ERL/00/000000/mmedia), were downloaded from KNMI Climate Explorer website database, <https://climexp.knmi.nl> (Trouet and van Oldenborgh 2013). A multi-model ensemble is used to account for uncertainty in the projections of the climate extremes, as both the ensemble mean, and associated spread, provide a more robust assessment of signal and noise than results from one or a few models (Tebaldi and Knutti 2007, Knutti *et al* 2010, IPCC 2012).

The indices are available re-gridded to a common spatial resolution of $2.5^\circ \times 2.5^\circ$ from their native model resolutions (Sillmann *et al* 2013b). Historical simulations (1861–2005) combined with the high emissions scenario Representative Concentration Pathway that projects an 8.5 W m^{-2} radiative forcing on the climate system by 2100 (RCP8.5) (Taylor *et al* 2012) are chosen for analysis. This RCP is chosen as the forcing is sufficiently intense to guarantee all models reach a warming of 2.0°C by the end of their simulations; further it is most representative of current forcing trends (Lewis and King 2017). As described below, we analyze indices from individual models at the time they reach specific global temperature targets, so the forcing scenario is not particularly important. Additionally, Pendergrass *et al* (2015) and Shi *et al* (2017) have shown that, in general, changes in climate extremes are indistinguishable across different RCPs when using multi-model ensembles; although Wang *et al* (2017) shows that differences in regional aerosol emissions do produce differences in projected changes over high emission areas (not Southern Africa).

This study focuses on extreme indices that relate directly to climate change vulnerability in Botswana (table 1) as determined from a review of vulnerability to climate in semi-arid countries of Southern Africa

Table 1. Temperature and precipitation climate extreme indices relevant to vulnerability assessment in Botswana. The indices are available from the KNMI Climate Explorer website [<http://climexp.knmi.nl>].

Symbol	Name	Units
Precipitation indices		
PRCPTOT	Annual total precipitation in wet days	mm/yr
ALTCDD	Maximum number of consecutive days per year with less than 1 mm of precipitation	days
ALTCWD	Maximum number of consecutive days per year with at least 1 mm of precipitation	days
RX1DAY	Annual maximum 1 day precipitation	mm/d
RX5DAY	Annual maximum 5 day precipitation	mm/5 d
R99P	Annual total precipitation when daily precipitation exceeds the 99th percentile of wet day precipitation	mm/yr
R95P	Annual total precipitation when daily precipitation exceeds the 95th percentile of wet day precipitation	mm/yr
R20MM	Annual count of days with at least 20 mm of precipitation	days
R10MM	Annual count of days with at least 10 mm of precipitation	days
Temperature indices		
TX90P	Percentage of days when daily maximum temperature is above the 90th percentile	%
TN90P	Percentage of days when daily minimum temperature is above the 90th percentile	%
TX10P	Percentage of days when daily minimum temperature is below the 10th percentile	%
TN10P	Percentage of days when daily minimum temperature is below the 10th percentile	%
WSDI	Maximum number of consecutive days per year when daily maximum temperature is above the 90th percentile	days

(Spear *et al* 2015). Rainfall deficit, rainfall variability and high temperature extremes have been found to be the main climate factors driving vulnerability (Batisani and Yarnal 2010, Omari 2010, Kgosikoma and Batisani 2014, Masundire *et al* 2016). PRCPTOT relates to annual rainfall deficits, R99P, R95P, RX1DAY, RX5DAY, R20MM and R10MM, ALTCWD heavy rainfall extremes and ALTCDD dry periods (irrespective of season). We note that rainfall extreme indices from GCMs may be biased compared to observations due to limitations in how models represent processes that drive precipitation extremes (Dai 2006, Westra *et al* 2014); they tend to precipitate more frequently, and in smaller amounts, affecting both dry spell length, as well the absolute value of heavy rainfall. However, within their own climate, they do simulate precipitation extremes (O’Gorman 2015), and so trends in these extremes can provide insights for the real world. For temperature-derived indices, WSDI is used to relate the potential impact of continuous high temperature instances (heat waves as defined by Moses 2017). TN90P, TN10P, TX10P and TN10P are also included to help in defining the potential impact of individual hot and cold events (Klein Tank *et al* 2009).

2.3. Methodology

A 40 year period (1861–1900) base period for the preindustrial era was defined from which the changes in extreme indices are compared following Huang *et al* (2017). The years at which each participating model reaches 1.0, 1.5 and 2.0 °C global warming above preindustrial levels is defined using a time sampling method initially used by Kaplan and New (2006) and applied by Zaroug *et al* (2018). The 1.0 °C temperature warming above preindustrial is used to represent the current climate (warming to date, from preindustrial) given the GMST reached 1.1 °C in 2016 (WMO 2017). A 31 year running mean is applied to the entire time-series for each model ensemble member. The

climatology at a given GMST warming above preindustrial is defined by the year the running mean reaches the GMST of interest and then stays consistently above the GMST. Figure 2 shows the spread of the years the participating models reach 1.0, 1.5 and 2.0 °C warming above preindustrial levels.


Monthly means of observed climate for 1979–2013 from the WATCH Forcing Data methodology applied to ERA-Interim data (WFDEI) (Weedon *et al* 2014) was used to cluster Botswana into three regions of homogeneous rainfall (figure 1(b)). The WFDEI dataset was found to simulate precipitation well over southern Africa (Li *et al* 2013) and has been used in a number of studies in sub-Saharan Africa (Andersson *et al* 2017, Nkiaka *et al* 2017). Shapefiles of Botswana and the three regions were created using ArcMap and used to extract the gridded sub-sets of the indices over each region, and the whole country.

Area-weighted average climatological means of the indices at a given GMST warming above pre-industrial levels were calculated within the subsets and used to determine the change relative to pre-industrial levels. For all the 24 members, the change for each extreme index relative to preindustrial levels is calculated as

$$\Delta I = I_n - I_0 \quad (1)$$

where I_n , $n \in \{1.0, 1.5 \text{ and } 2.0\}$ represents the area-averaged climatological mean calculated from the 31 year period surrounding the date of GMST, and I_0 is the area averaged climatological mean of the index of interest for 1861–1900 preindustrial times. Box-and-whisker plots of the absolute changes for each climate extreme index are plotted spanning the 24 member ensemble for each for the three regions and over the country areal average.

The non-parametric Wilcoxon Paired Signed Rank test (WPSR) is applied to test if the distributions of ensembles of the indices at 1.0, 1.5 and 2.0 °C are statistically significantly different from preindustrial levels,



and from each other. This test has been used in previous climate studies looking to determine the significance of changes in various climate indices/variables at different warming levels (Kharin *et al* 2013, Sillmann *et al* 2013b). To determine whether the models agree on the sign of change, a criterion that at least 75% of the members need to agree on the sign was adopted (Sillmann *et al* 2013b, Pinto *et al* 2016).

3. Results

Results are presented in box-and-whisker plot format, representing the ensemble spread for the change in the indices relative to the preindustrial baseline period. Changes are presented first for precipitation extreme indices followed by temperature extreme indices. From the plots, the ensemble median, interquartile range (IQR) and the outliers are represented. Ensemble member values that exceed $1.5 \times \text{IOR}$ are considered to be outliers. The box-and-whisker plots are made for each of the three regions (figure 1(b)) and the country average. Results obtained from testing model agreement on the sign of change are summarized in table S2 while table S3 summarizes the median and IQR changes in both of the precipitation and temperature indices relative to preindustrial levels. The inter-model ensemble spread, shown by the box-and-whisker plots together with the test for model agreement on change of sign depict the uncertainty in the projected changes of the indices. WPSR test results are presented using p -values.

3.1. Precipitation extremes changes

Changes in the precipitation indices at 1.0, 1.5 and 2.0 °C GMST warming above preindustrial levels indicate a progressive drying across Botswana, accompanied by an increase in heavy precipitation events, reduced wet spell events and increased dry spells (figure 3). The country average ensemble median change for total annual precipitation, PRCPTOT at 1.0, 1.5 and 2.0 °C GMST warming above preindustrial

levels indicate a reduction of -13 [range of -175 ; $+75$], -33 [-187 ; $+46$] and -63 [-192 ; $+40$] mm/yr (figure 3(a)). Of the three regions, Region 1 representing the northern and wettest parts of the country has the largest median reduction, and range, across the three warming periods with reductions of 20.2, 55.1 and 75.9 mm yr⁻¹ respectively (these are also the largest relative reductions, compared to preindustrial mean precipitation; see figure S1(b)). This region, as opposed to the rest of the country tends to be the one that receives most of its rainfall when the ITCZ shifts south in austral summer. The stabilizing and equatorward shifting of the mean position of the ITCZ under climate change could therefore be the reason behind the reduction (Giannini *et al* 2008). Region 3 has the least reductions (10–43 mm) in total annual precipitation, because of its already low annual precipitation totals (figure 1(a)); relative reductions are much larger (figure S1(a)). When testing for model agreement on change of sign, the change in PRCPTOT at 2.0 °C is the only one that shows consensus among models with $>75\%$ of the ensemble members projecting a reduction across all regions. Taking the country average, 58.3% of the models project a decrease in PRCPTOT at 1.0 °C, 62.5% at 1.5 °C and 83.3% at 2.0 °C GMST above preindustrial levels (table S2). WPSR test results show that the change in PRCPTOT between the three levels of warming is statistically significant across all regions, except for 1.0 versus 1.5 for a couple of the regions. An additional 0.5 °C increment in GMST from 1.5 °C and 2.0 °C therefore has significant impacts on the annual precipitation across the country. We note that PRCPTOT is not strictly a climate extreme, but it is included in climate extreme studies as total annual precipitation is a measure of interannual drought, and has implications on various economic sectors especially in water stressed countries like Botswana.

The number of consecutive dry days (ALTCDD) (figure 3(b)) show statistically significant increases across all regions for all warming levels. The ALTCDD median increases of 7.2, 13.5 and 18.7 days for the 1.0,





Figure 3. Box and whisker plots showing the changes in extreme precipitation indices across an ensemble of 24 participating model members. The changes are at 1.0, 1.5 and 2.0 °C above preindustrial levels (1861–1900). P-values from the WRS test are shown: P₀, P₁ and P₂ compares the ensemble spread of 1.0 °C–1.5 °C, 1.5 °C–2.0 °C and 1.0 °C–2.0 °C warmer climate regimes respectively.

1.5 and 2.0 °C respectively are projected for Region 1, these being the largest of the changes. On average, Botswana is projected to experience median increases of 5.7, 10.4 and 17.8 days in ALTCDD under the respective warmings. There is general consensus on the sign of change of ALTCDD across ensemble members with more than 80% of members depicting an increase across the three warming periods over the entire country. The increases in ALTCDD imply longer dry seasons with late onsets and early cessation of rainfall, as noted by Pinto *et al* (2016) and Sillmann *et al* (2013b). Median changes in the number of consecutive wet days (ALTCWD) are generally small in magnitude (figure 3(c)). Relative to preindustrial levels, median changes at 1.0 °C warming are for a decrease of

0.21 days in Region 1 while Region 2 and 3 show median increases of even smaller magnitude. On average, an additional 0.5 °C warming to 1.5 and 2.0 °C reduces ALTCWD by about half a day. The reductions in ALTCWD may be small in magnitude but are very significant given the short-lived and convective nature of rain bearing weather systems in Botswana. The shorter rainfall seasons described by increases in ALTCDD coupled with potentially shorter wet-spells could have serious implications on various economic sectors, with the agricultural sector likely to be particularly vulnerable.

For the heavy precipitation indices, the projected ensemble median changes in total accumulated precipitation from heavy (R95P) and very heavy (R99P)



precipitation days are a small, but generally non-significant increase as the climate system warms further (figures 3(d) and (e)). For R99P, the changes are mostly significant between the three warming levels in Regions 1 and 2, but not in Region 3. The ensemble spread generally disagrees on the sign of change for R95P, but there is more agreement for R99P, especially at the 1.5 °C and 2.0 °C warming levels.

Median changes in one-day and five-day maximum precipitation (RX1DAY and RX5DAY) show a general increase across all regions as the models progress to warmer climates. These changes are generally statistically significant over the wetter and semi-arid Regions 1 and 2, but not in the arid Region 3. (figures 3(f) and (g)). These results slightly contradict the findings by Sillmann *et al* (2013b) who concluded that there is a general decrease (though statistically insignificant) in RX5DAY in Southern Africa when looking at the changes for the period 2081–2100 relative to 1981–2000 Pinto *et al* (2016)'s findings using downscaled projections over Southern Africa are consistent with our results, though they looked the changes at 2069–2098 relative to the 1976–2005 period. Changes in the frequency of very heavy (R20MM) rainfall events do not show any significant changes, while the frequency of moderately heavy rainfall events (R10MM) show statistically significant decreases across all regions (figures 3(h) and (i)).

To investigate whether biases inherent in climate models especially in simulating accumulated precipita-

tion may influence the results, box-and-whisker plots of percentage changes in the total annual precipitation (PRCPTOT), one day maximum precipitation (RX1day) and the five-day maximum accumulated precipitation (RX5DAY) relative to industrial levels were also analysed (figure S1). Similar results to those obtained using absolute changes were found. An exception was that P_0 for PRCPTOT in Region 2 decreases from 0.117 when using absolute changes to 0.097 when using percentage changes (figure S1(a)). The difference suggests that the change in PRCPTOT between the current climate and 1.5 °C are statistically significantly different in Region 2. This could be because of the aridity in Region 2 meaning small percentage changes in total annual precipitation make a significant difference.

3.2. Temperature changes

The changes in temperature indices agree strongly on the direction of change across all indices (figure 4 and table S2). P -values for temperature derived indices obtained from the WPSR test are all very small ($\ll 0.10$) across all regions and warming levels above preindustrial levels. The p -values here imply that the changes in these indices are statistically significant across all warming levels. The inter-model spread is small for most temperature extremes therefore results are generally associated with less uncertainty.

Of the percentile based indices, the hot day and hot night extremes, TX90P (figure 4(a)) and TN90P

(figure 4(b)) show the greatest changes, though the change is more pronounced for TN90P especially over Region 1. TX90P is projected to increase by 30% above preindustrial levels on average in Botswana when the climate system reaches 2.0 °C, an increment of 10% from 1.5 °C levels. For TN90P the average increase at 2.0 °C is even higher, at 35%. Decreases in cold day and night extremes, TX10P (figure 4(c)) and TN10P (figure 4(d)), occur over the entire country with minimum temperature based extremes showing the larger reductions. Hot nights and mild winters are therefore expected to become a common occurrence with a warming climate leading to a decrease in frost occurrences. Warm spells (heat waves; WSDI) increase are projected across all regions, by 80 days compared to preindustrial levels at 2.0 °C for Region 1, and by 65 and 62 days for Region 2 and 3 (figure 4(e)). Even though all models show an increase in warm spells with increased warming, the ensemble spread also increases significantly at 2.0 °C (ranging between 26 and 96 days) compared to present (5–39 days) and 1.5 °C (13 and 65 days) indicating an increasing uncertainty as models warm further. Relating to findings by Moses (2017), the increases in TX90P and WSDI suggest a significant increase in heat wave events across the country. We note here the need to look at these indices seasonally as an opportunity for further investigation (Sillmann *et al* 2013b), and that the WSDI does not consider the intensity of heat waves (Dosio 2016).

3.3. Implications for vulnerability to climate change in Botswana

The projected changes in both temperature and precipitation extremes under warming climates are likely to have significant negative impacts on many social and economic activities in Botswana, most especially in the agricultural sector, and those dependent on water. The majority of Botswana's population is highly reliant on rain-fed agriculture for livelihoods, so these changes in extremes are likely to produce severe impacts, especially on the most vulnerable, women (Omari 2010). Many crops suffer sharp drops in yield after periods of cumulative heat stress (e.g. Schlenker and Lobell *et al* 2010). The large increases in temperature extremes projected for Botswana as one moves from 1.5 °C–2.0 °C, suggest potentially large, and growing impacts on crops that are currently farmed, such as maize (Barnabás *et al* 2008), significantly reducing yields. This is in agreement with previous studies showing that Botswana will be among the most impacted countries with regards to agriculture in Africa (Chipanshi *et al* 2003, Schlenker and Lobell 2010). These impacts will be exacerbated, particularly for rain-fed agriculture, by the projected decreases in mean annual rainfall and longer dry spells causing plant water stress, more frequently crossing the tipping point between good and average harvests, and complete agricultural failure (Batisani and Yarnal 2010). More intense rainfall will potentially cause crop damage and lower soil moisture.

These multiple impacts warn of a growing adaptation challenge as the climate warms from today, through 1.5–2.0 degrees, requiring more heat and drought-tolerant varieties to be developed or adopted. The options for expanding irrigated agriculture are small, apart from perhaps the North West of the country in the Okavango basin, but this would involve trade-offs with biodiversity conservation and ecotourism in this unique wetland system. Livestock production is also likely to be negatively impacted as increased dry spells will reduce pasture productivity (Setshwaelo 2001, Mberego 2017), compounded by inadequate adaptation strategies (Kgosikoma and Batisani 2014).

Water resources are also likely to be heavily impacted by the reduced total precipitation and increased intensity and longer dry spells, and greater evaporation under more extreme temperatures. Water stress is already a challenge with various economic sectors competing for the scarce resource; drought in 2014–2016 led to water-supply failure in the capital city, Gaborone (Siderius *et al* 2018). For Region 3, in the south-western parts of the country where rainfall is already low (annual accumulations of less than 300 mm), further reductions in rainfall imply increased pressure on the already stressed water sources (Batisani 2011). The increasing RX5DAY implies a potential increase in very heavy rainfall events that could cause flooding and lead to economic losses (Tsheko 2003). Though this is the case, some of these heavy rainfall events could come as a relief, replenishing stressed water resources. An example of such a case is the heavy downpours that came with post cyclone Dineo in February 2017, filling up the Gaborone dam that had run dry the previous year.

Climate sensitive diseases such as malaria will also be affected by the changes in climate extremes. As malaria epidemics thrive in wet and warmer climates, a general drying of the climate and shorter rainy seasons could lead to a reduction in the extent of the disease (Tanser *et al* 2003). Tanser *et al* (2003) also note that though for drier climates climate change might lead to a reduction in malaria incidences, epidemics can rise during times of heavy precipitation in these generally dry climates (Huang *et al* 2017). Based on the reasoning above, epidemics of malaria cases could also become a challenge as the climate systems warms to 2.0 °C. Other implications for health include increased chances of heat related mortalities as heat waves become a common occurrence, malnutrition due to reduced food supply as the agricultural sector is negatively impacted, and direct mortalities and injuries from floods (McMichael *et al* 2006).

4. Conclusions

This study found that Botswana is projected to experience significant increases in all temperature and many rainfall extremes as GMST increases from

1.0 °C through 1.5 °C–2.0 °C above preindustrial levels. The changes are particularly strong for temperature extremes; for each increment of global warming, temperature extremes are statistically different, indicating markedly different regional climates over Botswana at different levels of global warming. Similarly, there are projected statistically significant decreases in mean rainfall and increases in dry-spell length at each global temperature level. In contrast, although intense rainfall indices do show ensemble median increases, the spread in model results means that changes at each increment are not statistically distinguishable.

These changes in extremes present a growing adaptation challenge between 1.5 °C and 2.0 °C for key economic sectors in the country. Rain-fed agriculture is already marginal across much of the country, and the combined changes in heat extremes and decrease in moisture may well make current agricultural practices unviable at 1.5 °C and 2.0 °C warming. Botswana is already water-stressed; the projected decreases in mean annual rainfall, as well as increased dry spell length, will escalate stress, leading to more frequent water shortages in today's urban and agricultural supply systems. Further work is needed to better quantify the impacts, and resultant costs of adaptation at these different levels of global mean warming. However, our results suggest that, for a climate-stressed country such as Botswana, even small increments in global mean temperature have serious societal consequences that will demand progressively more radical adaptation responses.

A key limitation in our study relates to the use of GCM data, which are known to not completely represent extremes, especially rainfall, due to their relatively coarse spatial resolution. Further work is needed to apply downscaling methods to the GCM data, to add more information at finer space scales, subject to suitable evaluation of the downscaled data, as suggested by Pinto *et al* (2016). Another limitation to this study is that all extremes analysed are calculated on annual timescales; given that some of the indices such as ALTCDD and WSDI are also significant when looked at within seasons, further work is needed to look at the changes at seasonal timescales (Sillmann *et al* 2013b, Sillmann *et al* 2017).

Acknowledgments

This work was carried out under the Adaptation at Scale in Semi-Arid Regions project (ASSAR). ASSAR is one of four research programs funded under the Collaborative Adaptation Research Initiative in Africa and Asia (CARIAS), with financial support from the UK Government's Department for International Development (DfID) and the International Development Research Centre (IDRC), Canada. The views expressed in this work are those of the author and do not necessarily represent those of DfID and IDRC or its Board of Governors.

ORCID iDs

Tiro Nkemelang  <https://orcid.org/0000-0002-9588-7382>

Mark New  <https://orcid.org/0000-0001-6082-8879>

References

- Adams S *et al* 2013 *Turn Down the Heat: Climate Extremes, Regional Impacts, and the Case for Resilience* (Washington, DC: World Bank) (<http://documents.worldbank.org/curated/en/975911468163736818/Turn-down-the-heat-climate-extremes-regional-impacts-and-the-case-for-resilience-full-report>)
- Akinyemi F O 2017 Climate change and variability in semiarid palapye, eastern Botswana: an assessment from smallholder farmers' perspective *Weather Clim. Soc.* **9** 349–65
- Alexander L V *et al* 2006 Global observed changes in daily climate extremes of temperature and precipitation *J. Geophys. Res. Atmos.* **111** 1–22
- Andersson J C M, Ali A, Arheimer B, Gustafsson D and Minoungou B 2017 Providing peak river flow statistics and forecasting in the Niger River basin *Phys. Chem. Earth* **100** 3–12
- Andringa J 1984 The climate of Botswana in histograms *Botswana Notes Rec.* **16** 117–25
- ASC 2016 *UK Climate Change Risk Assessment 2017 Synthesis Report: Priorities for the Next Five Years* (London: Adaptation Sub-Committee of the Committee on Climate Change)
- Barnabás B, Jäger K and Fehér A 2008 The effect of drought and heat stress on reproductive processes in cereals *Plant Cell Environ.* **31** 11–38
- Batisani N 2011 Spatio-temporal ephemeral streamflow as influenced by climate variability in Botswana *J. Geog. Sci.* **21** 417–28
- Batisani N and Yarnal B 2010 Rainfall variability and trends in semi-arid Botswana: implications for climate change adaptation policy *Appl. Geogr.* **30** 483–9
- Bhalotra Y P R 1987 *Climate of Botswana, Part II: Elements of Climate* (Gaborone: Botswana Meteorological Services)
- Blamey R C and Reason C J C 2012 Mesoscale convective complexes over Southern Africa *J. Clim.* **25** 753–66
- Boko M *et al* 2008 *Africa Climate Change 2007: Impacts, Adaptation and Vulnerability. Contribution of Working Group II to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change* ed M L Parry, O Canziani, J Palutikof, P J van de Linden and C E Hanson (Cambridge: Cambridge University Press) pp 433–67
- Ceccherini G, Russo S, Ameztoy I, Francesco Marchese A and Carmona-Moreno C 2017 Heat waves in Africa 1981–2015, observations and reanalysis *Nat. Hazards Earth Syst. Sci.* **17** 115–25
- Chipanshi A C, Chanda R and Totolo O 2003 Vulnerability assessment of the maize and sorghum crops to climate change in Botswana *Clim. Change* **61** 339–60
- Cook C, Reason C J C and Hewitson B C 2004 Wet and dry spells within particularly wet and dry summers in the South African summer rainfall region *Clim. Res.* **26** 17–31
- Crimp S J and Mason S J 1999 The extreme precipitation event of 11–16 February 1996 over South Africa *Meteorol. Atmos. Phys.* **70** 29–42
- Dai A 2006 Precipitation characteristics in eighteen coupled climate models *J. Clim.* **19** 4605–30
- Dosio A 2016 Projection of temperature and heat waves for Africa with an ensemble of CORDEX regional climate models *Clim. Dyn.* **49** 493–519
- Driver P and Reason C J C 2017 Variability in the Botswana High and its relationships with rainfall and temperature characteristics over southern Africa *Int. J. Climatol.* **37** 570–81
- Favre A, Hewitson B, Lennard C, Cerezo-Mota R and Tadross M 2013 Cut-off lows in the South Africa region and their contribution to precipitation *Clim. Dyn.* **41** 2331–51

- Fischer E M, Beyerle U and Knutti R 2013 Robust spatially aggregated projections of climate extremes *Nat. Clim. Change* **3** 1033–8
- Frich P, Alexander L V, Della-Marta P, Gleason B, Haylock M, Tank Klein A M G and Peterson T 2002 Observed coherent changes in climatic extremes during the second half of the twentieth century *Clim. Res.* **19** 193–212
- Giannini A, Held I M and Sobel A H 2008 A global perspective on African climate *Clim. Change* **90** 359–83
- Hambira W L and Saarinen J 2015 Policy-makers' perceptions of the tourism–climate change nexus: policy needs and constraints in Botswana *Dev. South. Afr.* **32** 350–62
- Henley B J and King A D 2017 Trajectories towards the 1.5 °C Paris target: modulation by the Interdecadal Pacific Oscillation *Geophys. Res. Lett.* **44** 4256–62
- Huang J, Yu H, Dai A, Wei Y and Kang L 2017 Drylands face potential threat under 2 °C global warming target *Nat. Clim. Change* **7** 417–22
- IPCC 2012 *Managing the Risks of Extreme Events and Disasters to Advance Climate Change Adaptation* ed C B Field *et al* (Cambridge: Cambridge University Press)
- James R, Washington R, Schlessner C-F, Rogelj J and Conway D 2017 Characterizing half-a-degree difference: a review of methods for identifying regional climate responses to global warming targets *Wiley Interdiscip. Rev. Clim. Change* **8** e457
- Kaplan J O and New M 2006 Arctic climate change with a 2 °C global warming: Timing, climate patterns and vegetation change *Clim. Change* **79** 213–41
- Kgosikoma O E and Batisani N 2014 Livestock population dynamics and pastoral communities' adaptation to rainfall variability in communal lands of Kgalegadi South, Botswana *Pastoralism* **4** 19
- Kharin V V, Zwiers F W, Zhang X and Hegerl G C 2007 Changes in temperature and precipitation extremes in the IPCC ensemble of global coupled model simulations *J. Clim.* **20** 1419–44
- Kharin V V, Zwiers F W, Zhang X and Wehner M 2013 Changes in temperature and precipitation extremes in the CMIP5 ensemble *Clim. Change* **119** 345–57
- King A D, Karoly D J, Henley B J, Karoly D J and Henley B J 2017 Australian climate extremes at 1.5 °C and 2 °C of global warming *Nat. Clim. Change* **7** 412–16
- King D, Schrag D, Dadi Z, Ye Q and Ghosh A 2015 *Climate Change: A Risk Assessment* (Cambridge: University of Cambridge, Centre for Science and Policy) (www.csap.cam.ac.uk/projects/climate-change-risk-assessment)
- Klein Tank A M G, Zwiers F W and Zhang X 2009 *Guidelines on Analysis of Extremes in a Changing Climate in Support of Informed Decisions for Adaptation* (Geneva: World Meteorological Organization)
- Knutti R, Furrer R, Tebaldi C, Cermak J and Meehl G A 2010 Challenges in combining projections from multiple climate models *J. Clim.* **23** 2739–58
- Lewis S C and King A D 2017 Evolution of mean, variance and extremes in 21st century temperatures *Weather Clim. Extremes* **15** 1–10
- Li L, Ngongondo C S, Xu C and Gong L 2013 Comparison of the global TRMM and WFD precipitation datasets in driving a large-scale hydrological model in southern Africa *Hydrol. Res.* **44** 770–88
- Manhique A J, Reason C J C, Silinto B, Zucula J, Raiva I, Congolo F and Mavume A F 2015 Extreme rainfall and floods in southern Africa in January 2013 and associated circulation patterns *Nat. Hazards* **77** 679–91
- Manton M J 2010 Trends in climate extremes affecting human settlements *Curr. Opin. Environ. Sustain.* **2** 151–5
- Mason S J and Jury M R 1997 Climatic variability and change over southern Africa: a reflection on underlying processes *Progr. Phys. Geogr.* **21** 23–50
- Masundire H, Morchain D, Raditloang N, Hegga S, Ziervogel G, Molefe C, Angula M and Omari K 2016 *Vulnerability and Risk Assessment in Botswana's Bobirwa Sub-District: Fostering People-Centred Adaptation to Climate Change* (Ottawa: CARIAA and IDRC)
- Mberego S 2017 Temporal patterns of precipitation and vegetation variability over Botswana during extreme dry and wet rainfall seasons *Int. J. Climatol.* **37** 2947–60
- McElroy M and Baker D J 2012 *Climate Extremes: Recent Trends with Implications for National Security* (Cambridge: Harvard University Center for the Environment) (http://environment.harvard.edu/sites/default/files/climate_extremes_report_2012-12-04.pdf)
- McMichael A J, Woodruff R E and Hales S 2006 Climate change and human health: present and future risks *Lancet* **367** 859–69
- Molekwa S 2013 Cut-off lows over South Africa and their contribution to the total rainfall of the Eastern Cape Province *Master's thesis* (Pretoria: University of Pretoria)
- Monier E and Gao X 2015 Climate change impacts on extreme events in the United States: an uncertainty analysis *Clim. Change* **131** 67–81
- Moses O 2017 Heat wave characteristics in the context of climate change over the past 50 years in Botswana *Botswana Notes Rec.* **49** 13–25
- Moyo E N and Nangombe S S 2015 Southern Africa's 2012–13 violent storms: role of climate change *Procedia IUTAM* **17** 69–78
- ND-GAIN 2016 *Rankings IND-GAIN Country Index* (<https://gain.nd.edu/our-work/country-index/rankings/>) (Accessed: 8 February 2018)
- New M *et al* 2006 Evidence of trends in daily climate extremes over southern and west Africa *J. Geophys. Res. Atmos.* **111** 1–11
- Niang I, Ruppel O C, Abdrabo M A, Essel A, Lennard C, Padgham J and Urquhart P 2014 *Africa Climate Change 2014: Impacts, Adaptation and Vulnerability—Contributions of the Working Group II to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change* (Cambridge: Cambridge University Press) pp 1199–265
- Nicholson S E, Leposo D and Grist J 2001 The relationship between El Niño and Drought over Botswana *J. Clim.* **14** 323–35
- Nkiaka E, Nawaz N and Lovett J 2017 Evaluating global reanalysis datasets as input for hydrological modelling in the Sudano-Sahel Region *Hydrology* **4** 13
- O'Gorman P A 2015 Precipitation extremes under climate change *Curr. Clim. Change Rep.* **1** 49–59
- Omari K 2010 *Gender and Climate Change: Botswana Case Study* (Cape Town: Heinrich Böll Foundation Southern Africa) pp 49
- Pendergrass A G, Lehner F, Sanderson B M and Xu Y 2015 Does extreme precipitation intensity depend on the emissions scenario? *Geophys. Res. Lett.* **42** 8767–74
- Pinto I, Lennard C, Tadross M, Hewitson B, Dosio A, Nikulin G, Panitz H J and Shongwe M E 2016 Evaluation and projections of extreme precipitation over southern Africa from two CORDEX models *Clim. Change* **135** 655–68
- Razavi T, Switzman H, Arain A and Coulibaly P 2016 Regional climate change trends and uncertainty analysis using extreme indices: a case study of Hamilton, Canada *Clim. Risk Manage.* **13** 43–63
- Reason C J C 2007 Tropical cyclone Dera, the unusual 2000/01 tropical cyclone season in the South West Indian Ocean and associated rainfall anomalies over Southern Africa *Meteorol. Atmos. Phys.* **97** 181–8
- Reason C J C and Keibel A 2004 Tropical cyclone Eline and its unusual penetration and impacts over the Southern African Mainland *Weather Forecast.* **19** 789–805
- Richard Y, Fauchereau N, Pocard I, Rouault M and Trzaska S 2001 20th century droughts in southern Africa: spatial and temporal variability, teleconnections with oceanic and atmospheric conditions *Int. J. Climatol.* **21** 873–85
- Rouault M and Richard Y 2005 Intensity and spatial extent of droughts in southern Africa *Geophys. Res. Lett.* **32** L15702
- Sanderson B M *et al* 2017 Community climate simulations to assess avoided impacts in 1.5 °C and 2.0 °C futures *Earth Syst. Dyn. Discuss.* **8** 827–47

- Schlenker W and Lobell D B 2010 Robust negative impacts of climate change on African agriculture *Environ. Res. Lett.* **5** 14010
- Setshwaelo L 2001 *A Study of Grasslands and Livestock Vulnerability and Adaptation to Climate Change in Botswana* (Gaborone: Government of Botswana)
- Shi C, Jiang Z-H, Chen W-L and Li L 2017 Changes in temperature extremes over China under 1.5 °C and 2 °C global warming targets *Adv. Clim. Change Res.* **in press**
- Shongwe M E, Van Oldenborgh G J, Van Den Hurk B J J M, De Boer B, Coelho C A S and Van Aalst M K 2009 Projected changes in mean and extreme precipitation in Africa under global warming. Part I: Southern Africa *J. Clim.* **22** 3819–37
- Siderius C, Gannon K E, Ndiyoi M, Opere A, Batisani N, Olago D, Pardoe J and Conway D 2018 Hydrological response and complex impact pathways of the 2015/2016 El Niño in eastern and southern Africa *Earth's Future* **6** 2–22
- Sillmann J, Kharin V V, Zhang X, Zwiers F W and Bronaugh D 2013a Climate extremes indices in the CMIP5 multimodel ensemble: Part 1. Model evaluation in the present climate *J. Geophys. Res. Atmos.* **118** 1716–33
- Sillmann J, Kharin V V, Zwiers F W, Zhang X and Bronaugh D 2013b Climate extremes indices in the CMIP5 multimodel ensemble: Part 2. Future climate projections *J. Geophys. Res. Atmos.* **118** 2473–93
- Sillmann J *et al* 2017 Understanding, modeling and predicting weather and climate extremes: challenges and opportunities *Weather Clim. Extremes* **18** 65–74
- Singleton A T and Reason C J C 2007 Variability in the characteristics of cut-off low pressure systems over subtropical southern Africa *Int. J. Climatol.* **27** 295–310
- Spear D, Baudoïn M, Hegga S, Zaroug M, Okeyo A and Haimbili E 2015 *Vulnerability and Adaptation to Climate Change in the Semi-Arid Regions of Southern Africa* (Ottawa: CARIAA and IDRC)
- Tanser F C, Sharp B and Le Sueur D 2003 Potential effect of climate change on malaria transmission in Africa *Lancet* **362** 1792–8
- Taylor K E, Stouffer R J and Meehl G A 2012 An overview of CMIP5 and the experiment design *Bull. Am. Meteorol. Soc.* **93** 485–98
- Tebaldi C and Knutti R 2007 The use of the multi-model ensemble in probabilistic climate projections *Phil. Trans. R. Soc. A Math. Phys. Eng. Sci.* **365** 2053–75
- Trenberth K E 2012 Framing the way to relate climate extremes to climate change *Clim. Change* **115** 283–90
- Trouet V and van Oldenborgh G J 2013 KNMI climate explorer: a web-based research tool for high-resolution paleoclimatology *Tree-Ring Res.* **69** 3–13
- Tsheko R 2003 Rainfall reliability, drought and flood vulnerability in Botswana *Water SA* **29** 389–92
- UNFCCC 2015 *Decision 1/CP.21. The Paris Agreement* FCCC/CP/2015/L.9/Rev.1
- Usman M T and Reason C J C 2004 Dry spell frequencies and their variability over southern Africa *Clim. Res.* **26** 199–211
- Vossen P 1990 Rainfall and agricultural production in Botswana *Afrika Focus* **6** 141–55
- Wang Z, Lin L, Zhang X, Zhang H, Liu L and Xu Y 2017 Scenario dependence of future changes in climate extremes under 1.5 °C and 2 °C global warming *Sci. Rep.* **7** 46432
- Weedon G P, Balsamo G, Bellouin N, Gomes S, Best M J and Viterbo P 2014 The WFDEI meteorological forcing data set: WATCH Forcing Data methodology applied to ERA-Interim reanalysis data *Water Resour. Res.* **50** 7505–14
- Westra S *et al* 2014 Future changes to the intensity and frequency of short-duration extreme rainfall *Rev. Geophys.* **52** 522–55
- Williams C J R, Kniveton D R and Layberry R 2007 Climatic and oceanic associations with daily rainfall extremes over southern Africa *Int. J. Climatol.* **27** 93–108
- WMO 2017 *WMO Statement on the Status of the Global Climate in 2016* (1189) WMO-No 1189 (Geneva: World Meteorological Services)
- You Q *et al* 2011 Changes in daily climate extremes in China and their connection to the large scale atmospheric circulation during 1961–2003 *Clim. Dyn.* **36** 2399–417
- Zaroug M, New M and Lennard C 2018 private communication
- Zhang X, Alexander L, Hegerl G C, Jones P, Tank A K, Peterson T C, Trewin B and Zwiers F W 2011 Indices for monitoring changes in extremes based on daily temperature and precipitation data *Wiley Interdiscip. Rev. Clim. Change* **2** 851–70